

# ALCAPHOR

## DRINKABLE SOLUTION IN A BOTTLE

### ALCAPHOR

#### COMPOSITION

Trometamol citrate.....68,675g

(of which trometamol 38,325g)

Sodium citrate anhydrous.....3,675g

Dipotassium citrate anhydrous.....9,050g

Excipients : benzoic acid, sodium cyclamate, caramel,  
purified water (to make up the quantity of).....250 ml

#### PHARMACEUTICAL GROUP

Drinkable solution in a 250ml bottle

Authorised medicine N° 3400931268949

#### PHARMACO THERAPEUTIC GROUP :

Urinary alkalizer

#### WHEN TO USE THIS PRODUCT

This medicine is used for alkalinizing urine and to avoid the formation of acid kidney stones.

#### CONTRA-INDICATIONS

This medicine MUST NEVER BE USED in the following cases :

- anuria or severe renal impairment
- alkalosis
- diarrhea, in particular in patients who have recently undergone digestive surgery

IF IN DOUBT IT IS NECESSARY TO ASK THE ADVICE OF YOUR DOCTOR OR PHARMACIST

#### PRECAUTIONS FOR USE

This treatment can only be administered under medical supervision. This normally consists of controlling urinary PH.

**INTERACTIONS WITH OTHER MEDICINE AND OTHER INTERACTIONS**

In order to prevent possible interactions with other medicine, you must inform your doctor or pharmacist of any other treatment you are taking. This medicine contains benzoic acid

**DOSAGE**

The treatment must be adapted to each individual case. In any case, strictly follow your doctors instructions. The normal dosage is 2 to 4 tablespoons a day.

**METHOD AND ROUTE OF ADMINISTRATION**

By mouth

ALCAPHOR can be taken pure or diluted in half a glass of water, to be drunk slowly during meals.

**WHAT TO DO IN THE CASE OF OVERDOSE**

Immediately contact your doctor or a poison control centre.

**UNDESIRABLE EFFECTS**

LIKE ALL ACTIVE PRODUCTS, THIS MEDICINE MAY; IN CERTAIN CASES, CAUSE SIDE EFFECTS, WHICH ARE MORE OR LESS SERIOUS.

Intestinal transit may be accelerated during the first few days of the treatment : an anti-diarrhea medicine can be useful for a few days.

INFORM your doctor or your pharmacist if you notice ANY SIDE EFFECTS not mentioned in this leaflet.

**STORAGE :**

DO NOT USE AFTER THE EXPIRY DATE SHOWN ON THE PACK.

Keep at room temperature.

REVISION DATE : 10/97